

**DEVELOPMENT OF THREE PV SOLAR ENERGY
FACILITIES (EUPHORBIA PV, HILLARDIA PV &
VERBENA PV) AND ASSOCIATED INFRASTRUCTURE
NEAR LICHTENBURG, NORTH WEST PROVINCE**

Avifauna Scoping Report

February 2022



Compiled by:

Pachnoda Consulting CC
Lukas Niemand Pr.Sci.Nat

PO Box 72847
Lynwood Ridge
Pretoria
0040

Prepared for:

Euphorbia PV (Pty) Ltd, Verbena PV
(Pty) Ltd and Hillardia PV (Pty) Ltd

101, Block A, West Quay Building
7 West Quay Road, Waterfront
Cape Town, 8000

EXECUTIVE SUMMARY

Pachnoda Consulting cc was requested by Euphorbia PV (Pty) Ltd, Hillardia PV (Pty) Ltd and Verbena PV (Pty) Ltd to compile an avifauna scoping report for the proposed construction of three photovoltaic (PV) solar energy facilities (known as the Euphorbia PV, Hillardia PV and Verbena PV facilities), each with a contracted capacity of up to 100 MW located on a site approximately 10 km north west of the town of Lichtenburg in the North West Province.

The objectives of the scoping phase of the project were to obtain a basic overview of the variation and general status of the avifaunal habitat types and expected bird species likely to be affected by the proposed project.

Three avifaunal habitat types were identified, ranging from open mixed dolomite grassland with bush clump mosaics, moist grassland on low-lying areas and artificial livestock watering points. Approximately 181 bird species are expected to occur on the study area, including 11 Red listed species (threatened and near threatened species).

The main potential impacts associated with the proposed PV solar facility are expected to be the following:

- The loss of habitat and subsequent displacement of bird species due to the ecological footprint required during construction.
- Direct interaction (collision trauma) by birds with the surface infrastructure (photovoltaic panels) caused by polarised light pollution and/or waterbirds colliding with the panels (as they are mistaken for waterbodies).
- Collision with associated infrastructure (mainly overhead powerlines and reticulation)¹.

The endangered Cape Vulture (*Gyps coprotheres*), critically endangered White-backed Vulture (*Gyps africanus*) and Lappet-faced Vulture (*Torgos tracheliotos*) could occur as regular foraging visitors on the proposed PV facilities (according to reporting rates obtained from the atlas project - SABAP2). Although these species are highly prone to powerline collisions, the placement of the PV panels may result in the displacement of foraging individuals from the area. However, the risk of collision is considered more important when vultures feed on a carcass in close proximity to a powerline.

Collision risks associated with overhead powerlines (e.g. OHL and the grid corridor) will be dealt with in a separate Basic Assessment report, and such impacts will not form part of this screening report.

¹ It is important to note that impacts associated with the proposed powerline infrastructure will be assessed as part of a standalone Basic Assessment process and not as part of the facility EIAs. This baseline report does however report on potential powerline associated impacts and collision prone species as the two (the facilities and the grid connection infrastructure) are co-dependant and should be viewed holistically.

In addition, a total of 41 collision-prone bird species have been recorded from the study area (*sensu* atlas data), of which nine species could potentially interact with the PV panels. The study sites are not located near any prominent wetland system or impoundment, and therefore the risk of waterbird collisions with the proposed infrastructure (PV panels) was considered to be low.

TABLE OF CONTENTS

EXECUTIVE SUMMARY	I
TABLE OF CONTENTS	III
LIST OF FIGURES.....	IV
LIST OF TABLES.....	IV
LIST OF APPENDICES.....	IV
DECLARATION OF INDEPENDENCE	V
1. INTRODUCTION.....	1
1.1 PROJECT DESCRIPTION.....	1
1.2 TERMS OF REFERENCE	2
2. METHODS & APPROACH.....	5
2.1 LITERATURE SURVEY AND DATABASE ACQUISITION.....	5
2.2 PRELIMINARY SENSITIVITY ANALYSIS	6
2.3 LIMITATIONS	7
3. PRELIMINARY RESULTS AND DESCRIPTION OF THE AFFECTED ENVIRONMENT.....	9
3.1 LOCALITY	9
3.2 REGIONAL VEGETATION DESCRIPTION	9
3.3 LAND COVER, LAND USE AND EXISTING INFRASTRUCTURE.....	10
3.4 CONSERVATION AREAS, PROTECTED AREAS AND IMPORTANT BIRD AREAS	11
3.5 IMPORTANT AVIFAUNAL HABITAT TYPES	12
3.6 SPECIES RICHNESS AND PREDICTED SUMMARY STATISTICS.....	16
3.7 BIRD SPECIES OF CONSERVATION CONCERN	17
3.8 PRELIMINARY AVIFAUNAL SENSITIVITY	20
3.9 OVERVIEW OF AVIAN IMPACTS AT SOLAR FACILITIES.....	21
3.9.1 Background to solar facilities and their impact on birds	21
3.9.2 Potential impacts of PV solar facilities on birds	23
3.10 POTENTIAL IMPACTS ASSOCIATED WITH THE EUPHORBIA PV, HILLARDIA PV AND VERBENA PV SOLAR ENERGY FACILITIES.....	23
3.10.1 Loss of habitat and displacement of birds	23
3.10.2 Interaction with overhead powerlines (grid connection)	Error!
Bookmark not defined.	
4.10.3 Collision trauma caused by photovoltaic panels (the "lake-effect").....	24
4.10.4 Creation of "new" avian habitat and bird pollution.....	25
3.11 COLLISION-PRONE BIRD SPECIES	27
4. PLAN OF STUDY FOR THE EIA PHASE.....	32
4.1 PROPOSED APPROACH AND METHODS.....	32
5. REFERENCES.....	33

LIST OF FIGURES

Figure 1: A satellite image illustrating the geographic position of the proposed PV facilities.	4
Figure 2: A satellite image illustrating the regional vegetation type corresponding to the study site. Vegetation type categories were defined by Mucina & Rutherford (2006).....	10
Figure 3: A map illustrating the land cover classes (Geoterrainimage, 2015) corresponding to the proposed study site.	11
Figure 4: A map illustrating the locality of conservation areas in close proximity to the proposed study site.	12
Figure 5: A preliminary habitat map illustrating the avifaunal habitat types on the study site (the habitat types are subject to change pending the outcome of detailed surveys).	14
Figure 6: A collage of images illustrating examples of avifaunal habitat types on the study site observed during the austral summer season (January 2021): (a - d) open mixed dolomite grassland and bush clump mosaics, (e - f) moist grassland located on low-lying areas and (g - h) and artificial livestock watering points....	15
Figure 7: A map illustrating the preliminary avifaunal sensitivity of the area based on habitat types supporting bird taxa of conservation concern and important ecological function.....	21

LIST OF TABLES

Table 1: A summary table of the total number of species, Red listed species (according to Taylor et al., 2015 and the IUCN, 2021), endemics and biome-restricted species (Marnewick et al., 2015) expected (<i>sensu</i> SABAP1 and SABAP2) to occur in the study site.....	16
Table 2: Expected biome-restricted species (Marnewick <i>et al</i> , 2015) likely to occur on the study site.....	17
Table 3: Bird species of conservation concern that could utilise the study site based on their historical distribution range and the presence of suitable habitat. Red list categories according to the IUCN (2022)* and Taylor et al. (2015)**.....	18
Table 4: Collision-prone bird species and Red listed species (in red) expected to be present on the study site.	29

LIST OF APPENDICES

Appendix 1: A shortlist of bird species expected to be present on the study area. The list provides an indication of the species occurrence according to SABAP2 reporting rates.....	36
--	----

DECLARATION OF INDEPENDENCE

I, Lukas Niemand (Pachnoda Consulting CC) declare that:

- I act as the independent specialist in this application to Euphorbia PV (Pty) Ltd, Hillardia PV (Pty) Ltd, Verbena PV (Pty) Ltd and Atlantic Renewable Energy Partners (Pty) Ltd;
- I will perform the work relating to the application in an objective manner, even if this results in views and findings that are not favourable to the applicant;
- I declare that there are no circumstances that may compromise my objectivity in performing such work;
- I have no vested financial, personal or any other interest in the application;
- I have no, and will not engage in, conflicting interests in the undertaking of the activity;
- I undertake to disclose to the applicant and the competent authority all material information in my possession that reasonably has or may have the potential of influencing - any decision to be taken with respect to the application by the competent authority; and - the objectivity of any report, plan or document to be prepared by myself for submission to the competent authority; and
- All the particulars furnished by me in this form are true and correct.



Lukas Niemand (Pr.Sci.Nat)

11 February 2022

Lukas Niemand is registered with The South African Council for Natural Scientific Professionals (400095/06) with more than 15 years of experience in ecological-related assessments and more than 15 years in the field of bird interactions with electrical and renewable energy infrastructure. He has conducted numerous ecological and avifaunal impact assessments including Eskom Transmission projects, hydro-electric schemes, solar farms and other activities in South Africa and other African countries.

1. INTRODUCTION

1.1 Project Description

Pachnoda Consulting cc was requested by Euphorbia PV (Pty) Ltd, Hillardia PV (Pty) Ltd and Verbena PV (Pty) Ltd to compile an avifauna scoping report for the proposed construction of three photovoltaic (PV) solar energy facilities (known as the Euphorbia PV, Hillardia PV and Verbena PV facilities), each with a contracted capacity of up to 100 MW located on a site approximately 10 km north west of the town of Lichtenburg in the North West Province (Figure 1). The development area is situated within the Ditsobotla Local Municipality within the Ngaka Modiri Molema District Municipality. The site is accessible via an existing gravel road which provides access to the development area off the R505, located east of the development area.

The infrastructure of each proposed facility will consist of the following components:

- PV modules and mounting structures;
- Inverters and transformers;
- Battery Energy Storage System (BESS);
- Site and internal access roads (up to 8m wide);
- Auxiliary buildings (22kV or 33kV switch room, gate-house and security, control centre, office, warehouse, canteen & visitors centre, staff lockers etc);
- Temporary and permanent laydown area;
- Cabling between the panels, to be laid underground where practical; and
- Grid connection solution, including:
 - Medium-voltage underground cabling between the project components and the facility substation (within a 100 m wide and 1.5 km in length corridor for the Verbena PV facility and 2.5 km for the Hillardia PV facility). The MV cabling corridor is only applicable to Hillardia and Verbena facilities only, as the facility substation for Euphorbia will be on-site); and
 - A 132kV facility substation.

1.1.1 Euphorbia PV Facility

The development area for the Euphorbia PV facility and associated infrastructure will be located on the following properties:

- Portion 2 of the Farm Houthaalboomen 31
- Portion 3 of the Farm Houthaalboomen 31
- Portion 4 of the Farm Houthaalboomen 31

The development area of the Euphorbia PV facility is approximately 207 ha.

1.1.2 Hillardia PV Facility

The development area for the Hillardia PV facility and associated infrastructure will be located on the following properties:

- Portion 2 of the Farm Houthaalboomen 31
- Portion 3 of the Farm Houthaalboomen 31
- Portion 4 of the Farm Houthaalboomen 31

The development area of the Hillardia PV facility is approximately 230 ha.

1.1.3 Verbena PV Facility

The development area for the Verbena PV facility and associated infrastructure will be located on the following properties:

- Portion 2 of the Farm Houthaalboomen 31
- Portion 3 of the Farm Houthaalboomen 31
- Portion 4 of the Farm Houthaalboomen 31

The development area of the Verbena PV facility is approximately 220 ha.

Each facility will include grid connection infrastructure (MV cabling and facility substation) that will facilitate the connection of the project components to the Houthaalboomen North collector switching station which will be located adjacent to the Euphorbia PV facility.

The Houthaalboomen North collector switching station intends to connect the Houthaalboomen North cluster to the National Grid via the Watershed Main Transmission Substation (MTS) (approximately 5 km southeast of the facility), however, the connection infrastructure associated with this grid solution (i.e. between the facility substations and the MTS) is being assessed as part of a separate Environmental Application..

1.2 Terms of Reference

The main aim of this scoping exercise was to investigate the avifaunal attributes of the proposed PV facilities by means of a desktop analysis of GIS based information and third-party datasets and included a brief site visit which constituted the austral summer season sampling survey (January 2022).

Since the three proposed PV facilities are spatially autocorrelated (located within the same broad-scale habitat types, topography and climatic conditions), a combined scoping report was compiled which aims to provide an overview of the avifaunal attributes on the following properties ("herewith referred to as the "study site"):

- Portion 2 of the Farm Houthaalboomen 31
- Portion 3 of the Farm Houthaalboomen 31
- Portion 4 of the Farm Houthaalboomen 31

However, a separate avifaunal impact report will be compiled for each facility (three reports) during the EIA phase of the project.

The terms of reference for this scoping report are to:

- conduct an assessment on a screening level based on available information pertinent to the ecological and avifaunal attributes on the study site;
- conduct an assessment of all information on a screening level in order to present the following results:
 - typify the regional vegetation and avifaunal macro-habitat parameters that will be affected by the proposed project;
 - provide an indication on the occurrence of threatened, near-threatened, endemic and conservation important bird species likely to be affected by the proposed project;
 - provide an indication of sensitive areas or bird habitat types corresponding to the study site;
 - highlight areas of concern or "hotspot" areas;
 - identify potential impacts that are considered pertinent to the proposed development;
 - highlight gaps of information in terms of the avifaunal environment; and
 - recommend further studies to be conducted as part of the Environmental Impact Assessment (EIA) phase.




Figure 1: A satellite image illustrating the geographic position of the proposed PV facilities.

2. METHODS & APPROACH

The objectives of this phase of the project were to obtain a basic overview of the variation and general status of the avifaunal habitat types and expected bird species likely to be affected by the proposed project.

Also take note that the current report put emphasis on the avifaunal community as a key indicator group on the proposed study area, thereby aiming to describe the preliminary conservation significance of the ecosystems in the area. Therefore, the occurrence of certain bird species and their relative abundances (to be determined during the EIA although herewith deduced from SABAP2 reporting rates) could determine the outcome of the ecological sensitivity of the area and the subsequent layout of the proposed solar facility infrastructure.

The information provided in this report was principally sourced from the following sources/observations:

- relevant literature – see section below;
- observations made during a site visit (17 - 19 January 2022); and
- personal observations from similar habitat types in proximity to the study area, with emphasis on assessments conducted by Pachnoda Consulting (2018; 2021) of where an avifauna study was conducted by the author.

2.1 Literature survey and Database acquisition

A desktop and literature review of the area under investigation was commissioned to collate as much information as possible prior to the detailed baseline survey. Literature consulted primarily makes use of small-scale datasets that were collected by citizen scientists and are located at various governmental and academic institutions (e.g. Animal Demography Unit & SANBI). These include (although are not limited to) the following:

- Hockey *et al.* (2005) for general information on bird identification and life history attributes.
- Marnewick *et al.* (2015) was consulted for information regarding the biogeographic affinities of selected bird species that could be present on the study area.
- The conservation status of bird species was categorised according to the global IUCN Red List of threatened species (IUCN, 2022) and the regional conservation assessment of Taylor *et al.* (2015).
- Distributional data was sourced from the South African Bird Atlas Project (SABAP1) and verified against Harrison *et al.* (1997) for species corresponding to the quarter-degree grid cell (QDGC) 2626AA (Lichtenburg). The information was then modified according to the prevalent habitat types present on the study area. The SABAP1 data provides a “snapshot” of the abundance and composition of species recorded within a quarter degree grid cell (QDGC)

which was the sampling unit chosen (corresponding to an area of approximately 15 min latitude x 15 min longitude). It should be noted that the atlas data makes use of reporting rates that were calculated from observer cards submitted by the public as well as citizen scientists. It therefore provides an indication of the thoroughness of which the QDGCs were surveyed between 1987 and 1991.

- Additional distributional data was also sourced from the SABAP2 database (<http://www.sabap2.birdmap.africa>). The information was then modified according to the prevalent habitat types present on the study area. Since bird distributions are dynamic (based on landscape changes such as fragmentation and climate change), SABAP2 was born (and launched in 2007) from SABAP1 with the main difference being that all sampling is done at a finer scale known as pentad grids (5 min latitude x 5 min longitude, equating to 9 pentads within a QDGC). Therefore, the data is more site-specific, recent and more comparable with observations made during the site visit (due to increased standardisation of data collection). The pentad grids relevant to the current project is 2600_2600 and 2600_2605 (although all eight pentad grids surrounding the central grid 2600_2600 were also scrutinised with emphasis on grid 2605_2605).
- **The expected bird list was inferred from personal observations obtained during previous surveys (2021 and 2018) in the study area (from neighbouring farm portions on the Farm Houthaalboomen 31).**
- The choice of scientific nomenclature, taxonomy and common names were recommended by the International Ornithological Committee (the IOC World Bird List v. 12.1), unless otherwise specified (see www.worldbirdnames.org as specified by Gill et al, 2022).
- The incidental occurrence records for large birds of prey and vulture tracking data were included (only for 2018).
- Data on power line derived bird mortalities were requested from the electrical infrastructure mortality incident register (the dataset was provided by EWT).
- The best practice guidelines for assessing and monitoring the impact of solar power generating facilities on birds in southern Africa were also consulted (Jenkins *et al.*, 2017).
- Additional information regarding interactions of birds with PV facilities was provided by the author's own personal observations.

2.2 Preliminary Sensitivity Analysis

A preliminary sensitivity map was compiled based on the outcome of a desktop analysis.

The ecological sensitivity of any piece of land is based on its inherent ecosystem service (e.g. wetlands) and overall preservation of biodiversity.

2.3.1 Ecological Function

Ecological function relates to the degree of ecological connectivity between systems within a landscape matrix. Therefore, systems with a high degree of landscape connectivity amongst one another are perceived to be more sensitive and will be those contributing to ecosystem service (e.g. wetlands) or the overall preservation of biodiversity.

2.3.2 Avifaunal Importance

Avifaunal importance relates to species diversity, endemism (unique species or unique processes) and the high occurrence of threatened and protected species or ecosystems protected by legislation.

2.3.3 Sensitivity Scale

- *High* – Sensitive ecosystems with either low inherent resistance or low resilience towards disturbance factors or highly dynamic systems considered important for the maintenance of ecosystem integrity. Most of these systems represent ecosystems with high connectivity with other important ecological systems OR with high species diversity and usually provide suitable habitat for a number of threatened or rare species. These areas should preferably be protected;
- *Medium* – These are slightly modified systems which occur along gradients of disturbances of low-medium intensity with some degree of connectivity with other ecological systems OR ecosystems with intermediate levels of species diversity but may include potential ephemeral habitat for threatened species; and
- *Low* – Degraded and highly disturbed/transformed systems with little ecological function and are generally very poor in species diversity (most species are usually exotic or weeds).

2.3 Limitations

To obtain a comprehensive understanding of the diversity and dynamics of avifaunal community on the study area, as well as the status of endemic, rare or threatened species in the area, detailed assessments should always consider investigations at different time scales (across seasons/years) and through replication. However, due to the fact that the findings in this report were based on a scoping/screening assessment, long-term studies were not feasible and inferred interpretations were mostly based on ad hoc observations.

It should also be realised that bird distribution patterns fluctuate widely in response to environmental conditions (e.g. local rainfall patterns, nomadism, migration patterns, seasonality), meaning that a composition noted at a particular moment in time will differ

during another time period at the same locality. For this reason a dry season and wet season survey will be conducted.

Due to the scope of the work presented during a scoping assessment, a detailed investigation of the avifaunal community in the area were not possible and is not perceived as part of the Terms of Reference for a scoping/screening level exercise.

Furthermore, additional information may become known during a later stage of the process or development. This company, the consultants and/or specialist investigators do not accept any responsibility for conclusions, suggestions, limitations and recommendations made in good faith, based on the information presented to them, obtained from the surveys or requests made to them at the time of this report.

The following assumptions are relevant to the literature survey and database acquisition phase:

- It is assumed that third party information (obtained from government, academic/research institution, non-governmental organisations) is accurate and true;
- Some of the datasets are out of date and therefore extant distribution ranges may have shifted although these datasets could provide insight into historical distribution ranges of relevant species;
- The datasets are mainly small-scale and could not always consider azonal habitat types that may be present on the study area (e.g. small dams, pans and depressions). In addition, these datasets encompass surface areas larger than the study area that could include habitat types and species that is not present on the study area. Therefore, the potential to overestimate species richness is highly likely while it is also possible that certain cryptic or specialist species could have been overlooked in the past;
- Some of the datasets (e.g. SABAP2) managed by the Animal Demography Unit of the University of Cape Town were only recently initiated and therefore incomplete; and
- In addition, the study site is under private ownership and primarily inaccessible to the public. Since most of the species distribution ranges concerning the relevant datasets are subject to observations made by the public, it is likely that many bird species are overlooked or not formally catalogued for the area.

3. PRELIMINARY RESULTS AND DESCRIPTION OF THE AFFECTED ENVIRONMENT

3.1 Locality

The proposed PV facilities comprise of Portions 2, 3 and 4 of the Farm Houthaalboomen 31, located approximately 10 km north west of the town of Lichtenburg in the North West Province (Figure 1).

3.2 Regional Vegetation Description

The proposed PV facilities correspond to the Grassland Biome and more particularly to the Dry Highveld Grassland Bioregion as defined by Mucina & Rutherford (2006). It comprehends an ecological type known as Carletonville Dolomite Grassland (Mucina & Rutherford, 2006) (Figure 2).

From an avifaunal perspective it is evident that bird diversity is positively correlated with vegetation structure, and floristic richness is not often regarded to be a significant contributor of patterns in bird abundance and their spatial distributions. Although grasslands are generally poor in woody plant species, and subsequently support lower bird richness values, it is often considered as an important habitat for many terrestrial bird species such as larks, pipits, korhaans, cisticolas, widowbirds including large terrestrial birds such as Secretarybirds, cranes and storks. Many of these species are also endemic to South Africa and display particularly narrow distribution ranges. Due to the restricted spatial occurrence of the Grassland Biome and severe habitat transformation, many of the bird species that are restricted to the grasslands are also threatened or experiencing declining population sizes.

Carletonville Dolomite Grassland is confined to the dolomite plains that stretch from Lichtenburg in the North West Province to sections of rocky grassland in Gauteng, especially between altitudes of 1 350 m and 1 450 m. It occurs on slightly undulating plains dissected by prominent chert ridges, thereby containing a grassland composition rich in floristic species forming a complex mosaic dominated by many plant species.

Currently, only 2 % of the remaining 76 % of untransformed Carletonville Dolomite Grassland is formally protected within the Cradle of Humankind World Heritage Site and various nature reserves such as Abe Baily and Krugersdorp Nature Reserves.




Figure 2: A satellite image illustrating the regional vegetation type corresponding to the study site. Vegetation type categories were defined by Mucina & Rutherford (2006).

3.3 Land cover, land use and existing infrastructure.

According to the South African National dataset of 2013-2014 (Geoterrainimage, 2015) the study site comprehends the following land cover categories (Figure 3):

Natural areas:

- Grassland;
- Low shrubland; and
- Woodland and open bush.

From the land cover dataset it is evident that most of the study site is covered by natural grassland, while some parts consist of low shrubland, especially on the eastern (mainly Euphorbia PV facility) and south-western section of the study site (mainly Hillardia PV facility). The study site is primarily used for livestock production and livestock grazing. Existing infrastructure includes two cattle kraals/watering points located on the Euphorbia and Hillardia PV facilities.




Figure 3: A map illustrating the land cover classes (Geoterrainimage, 2015) corresponding to the proposed study site.

3.4 Conservation Areas, Protected Areas and Important Bird Areas

The study site is located approximately 3.2 km west of the former Lichtenburg Game Breeding Centre (Figure 4). This conservation area contains a variety of game species, and the facility operates a vulture restaurant which attracts foraging vultures (c. three species) to the region. This area is currently under new management (by lease agreement with the municipality).

There are no other formal protected areas or any Important Bird and Biodiversity Areas in close proximity to the study site.




Figure 4: A map illustrating the locality of conservation areas in close proximity to the proposed study site.

3.5 Important avifaunal habitat types

Apart from the regional vegetation type, the local composition and distribution of the vegetation associations on the study site are a consequence of a combination of factors simulated by soil type, geology and grazing intensity (presence of livestock) which have culminated in a number of habitat types that deserve further discussion² (Figure 5 and Figure 6):

1. *Open mixed dolomite grassland with bush clump mosaics:* This unit is prominent on the study site and covers a significant extent in surface area of the proposed PV facilities. It is represented by two discrete floristic variations which also provide habitat for two discrete avifaunal associations (see Pachnoda Consulting, 2018; 2021). The first floristic variation consists of open untransformed to grazed mixed dolomite grassland and bush clumps with an eminent woody layer. The grassland variation is represented by untransformed to grazed Carletonville Dolomite Grassland, depending on grazing intensity, and dominated by "late-successional" graminoids such a *Themeda triandra*, *Cymbopogon caesius*, *C. pospischilii*, *Trachypogon spicatus*, *Elionurus muticus* and *Andropogon schirensis*. It is occupied by a typical grassland bird

² The habitat types are subject to change pending on the outcome of a detailed austral summer survey.

composition dominated by insectivorous and granivore passerine bird species such as Desert Cisticola, (*Cisticola aridulus*), Eastern Clapper Lark (*Mirafra fasciolata*), Spike-heeled Lark (*Chersomanes albofasciata*), Ant-eating Chat (*Myrmecocichla formicivora*), Rufous-naped Lark (*Mirafra africana*) and Cloud Cisticola (*Cisticola textrix*). Prominent non-passerine species include Orange River Francolin (*Scleroptila gutturalis*), Swainson's Spurfowl (*Pternistis swainsonii*), Northern Black Korhaan (*Afrotis afraoides*), Crowned Lapwing (*Vanellus coronatus*) and Helmeted Guineafowl (*Numida meleagris*).

The bush clumps form a prominent mosaic characterised by the dominance of a woody layer of *Searsia lancea*, *S. pyroides*, *Senegalia cf. hereroensis* and *Vachellia erioloba* forms canopy constituents in some areas. The eminent increase in vertical heterogeneity provided by the woody layer is colonised by a "Bushveld" bird association consisting of insectivorous passerines such as Black-chested Prinia (*Prinia flavicans*), Chestnut-vented Warbler (*Curruca subcoerulea*), Kalahari Scrub Robin (*Cercotrichas paena*), Pycnonotus nigricans (*Cisticola fulvicapilla*), African Red-eyed Bulbul (*Pycnonotus nigricans*) as well as granivores such as Yellow Canary (*Crithagra flaviventris*), White-browed Sparrow-Weaver (*Plocepasser mahali*) and Southern Masked Weaver (*Ploceus velatus*). Non-passerine bird taxa are represented by Laughing Dove (*Spilopelia senegalensis*), Ring-necked Dove (*Streptopelia capicola*), Acacia Pied Barbet (*Tricholaema leucomelas*) and White-backed Mousebird (*Colius colius*).

2. *Moist grassland located within low-lying areas*: This habitat is located on the north-eastern part of the study site and receives infiltration from run-off water during precipitation events. It is colonised by dense, coarse grass including dense *Hyparrhenia*, *Themeda triandra* and *Heteropogon contortus* which provide breeding and roosting habitat for Long-tailed Widowbird (*Euplectes progne*), Southern Red Bishop (*E. orix*) and Zitting Cisticola (*Cisticola juncidis*). It is also often visited by terrestrial species such as Blacksmith Lapwing (*Vanellus armatus*).
3. *Artificial livestock watering points*: These are represented by artificial water troughs and reservoirs with the purpose to provide drinking water to livestock. It was only present on the proposed Hillardia and Euphorbia PV facilities. However, they act as focal congregation areas for many granivore passerine and non-passerine species, including Cape Sparrow (*Passer melanurus*), Laughing Dove (*Spilopelia senegalensis*), Namaqua dove (*Oena capensis*), Scaly-feathered Weaver (*Sporopipes squamifrons*) and Wattled Starling (*Creatophora cinerea*).
4. *Transformed areas*: These are represented by an *excavation* on the southern boundary of the proposed Verbena PV facility. It is an unimportant habitat for bird species.




Figure 5: A preliminary habitat map illustrating the avifaunal habitat types on the study site (the habitat types are subject to change pending the outcome of detailed surveys).






Figure 6: A collage of images illustrating examples of avifaunal habitat types on the study site observed during the austral summer season (January 2021): (a - d) open mixed dolomite grassland and bush clump mosaics, (e - f) moist grassland located on low-lying areas and (g - h) and artificial livestock watering points.

3.6 Species Richness and Predicted summary statistics

Approximately ~181 bird species are expected to occur on the study site and immediate surroundings (refer to Appendix 1 & Table 1). The expected richness was inferred from the South African Bird Atlas Project (SABAP1 & SABAP2) (Harrison et al., 1997; www.sabap2.birdmap.africa), personal observations and the presence of suitable habitat in the study area. The expected richness is also strongly correlated with favourable environmental conditions (e.g. during good rains) and seasonality (e.g. when migratory species are present). This equates to 18 % of the approximate 986³ species listed for the southern African subregion⁴ (and approximately 21 % of the 858 species recorded within South Africa⁵). However, the average species richness obtained from the pentad grid 2600_2600 and 2600_2605 corresponding to the study site contained 110 species, with an average number of 42 species for each full protocol card submitted (for observation of two hours or more). According to personal observations, the average number of species observed on the study site range between 60 - 75 bird species (obtained during the austral summer season of January 2021).

According to Table 1, the study site is expected to be poorly represented by biome-restricted (see Table 2) and local endemic bird species. It is expected to support ca. 34 % of the near-endemic species present in the subregion. Of the 181 bird species expected to occur in the project area, 11 are threatened or near threatened species, 15 are southern African endemics and 21 are near-endemic species. In addition, two threatened species (White-backed Vulture *Gyps africanus* and Cape Vulture *G. coprotheres*) were observed on the study site (Table 3).

Table 1: A summary table of the total number of species, Red listed species (according to Taylor et al., 2015 and the IUCN, 2021), endemics and biome-restricted species (Marnewick et al., 2015) expected (*sensu* SABAP1 and SABAP2) to occur in the study site.

Description	Expected Richness Value***
Total number of species*	181 (21 %)
Number of Red Listed species*	11 (8 %)
Number of biome-restricted species – Zambezi and Kalahari-Highveld Biomes)*	4 (29 %)
Number of local endemics (BirdLife SA, 2018)*	2 (5 %)
Number of local near-endemics (BirdLife SA, 2018)*	7 (23 %)
Number of regional endemics (Hockey <i>et al.</i> , 2005)**	15 (14 %)
Number of regional near-endemics (Hockey <i>et al.</i> , 2005)**	21 (34 %)

* only species in the geographic boundaries of South Africa (including Lesotho and Swaziland) were considered.

** only species in the geographic boundaries of southern Africa (including Namibia, Botswana, Zimbabwe and Mozambique south of the Zambezi River) were considered

3 *sensu* www.zestforbirds.co.za (Hardaker, 2020)

4 A geographical area south of the Cunene and Zambezi Rivers (includes Namibia, Botswana, Zimbabwe, southern Mozambique, South Africa, Swaziland and Lesotho).

5 With reference to South Africa (including Lesotho and Swaziland (BirdLife South Africa, 2018).

*** Percentage values in brackets refer to totals compared against the South African avifauna (*sensu* BirdLife SA, 2018).

Table 2: Expected biome-restricted species (Marnewick *et al*, 2015) likely to occur on the study site.

Species	Kalahari-Highveld	Zambezian	Expected Frequency of occurrence
Kalahari Scrub-robin (<i>Cercotrichas paena</i>)	X		Common
Kurichani Thrush (<i>Turdus libonyana</i>)		X	Uncommon
White-throated Robin-chat (<i>Cossypha humeralis</i>)		X	Rare
White-bellied Sunbird (<i>Cinnyris talatala</i>)		X	Uncommon

3.7 Bird species of conservation concern

Table 3 provides an overview of bird species of conservation concern that could occur on the study site based on their historical distribution ranges and the presence of suitable habitat. According to Table 3, a total of 11 species could occur on the study site which includes six globally threatened species, one globally near threatened species, two regionally threatened species and two regionally near-threatened species.

It is evident from Table 3 that the highest reporting rates (>5%) were observed for the globally endangered Cape Vulture (*Gyps coprotheres*) and the globally critically endangered White-backed Vulture (*Gyps africanus*). These species have a high likelihood of occurrence pending the presence of suitable food (livestock carcasses).

The regionally vulnerable Lanner Falcon (*Falco biarmicus*), globally endangered Lappet-faced Vulture (*Torgos tracheliotos*) and globally near threatened Red-footed Falcon (*Falco vespertinus*) show reporting rates between 2% and 5 %. These species have a moderate probability of occurrence and are regarded as occasional foraging visitors to the area.

The remaining species have low reporting rates (<2%) and are regarded as irregular foraging visitors with low probabilities of occurrence. However, during the brief scoping site visit it was noticed that extensive areas of suitable foraging habitat persists for some of these species (e.g. Secretarybird *Sagittarius serpentarius*) despite being ominously absent from the area. It is possible that the low reporting rates reflect the poor coverage of the study area by citizen scientists (e.g. birdwatchers), and some of these species could occur in higher numbers due to being overlooked. As an example, Red-footed Falcons (*F. vespertinus*) often occur in flocks of the similar-looking Amur Falcon (*F. amurensis*), which based on reporting rates appear to be a common summer visitor to the area. Therefore, it is highly possible that Red-footed Falcons were previously overlooked or misidentified.

Table 3: Bird species of conservation concern that could utilise the study site based on their historical distribution range and the presence of suitable habitat. Red list categories according to the IUCN (2022)* and Taylor et al. (2015).**

Species	Global Conservation Status*	National Conservation Status**	Mean Reporting rate: SABAP1	Mean Reporting rate: SABAP2	Preferred Habitat	Potential Likelihood of Occurrence
<i>Anthropoides paradiseus</i> (Blue Crane)	Vulnerable	Near threatened	47.18	-	Prefers open grasslands. Also forages in wetlands, pastures and agricultural land.	Potential vagrant or highly irregular foraging visitor.
<i>Aquila rapax</i> (Tawny Eagle)	Endangered-	Endangered	2.11	-	Lowveld and Kalahari savannas, especially game farming areas and reserves	An irregular visitor or vagrant to the study site.
<i>Ciconia abdimii</i> (Abdim's Stork)	-	Near threatened	7.75	-	Open stunted grassland, fallow land and agricultural fields.	An uncommon summer foraging visitor to areas consisting of secondary grassland or arable land.
<i>Falco vespertinus</i> (Red-footed Falcon)	Near threatened	Near threatened	2.11	3.13	Varied, prefers to hunt open arid grassland and savannoid woodland, often in company with Amur Falcons (<i>F. amurensis</i>).	An occasional summer foraging visitor to the area.
<i>Falco biarmicus</i> (Lanner Falcon)	-	Vulnerable	2.82	6.7 (for pentad 2600_2605)	Varied, but prefers to breed in mountainous areas.	An occasional foraging visitor to the study area.

Species	Global Conservation Status*	National Conservation Status**	Mean Reporting rate: SABAP1	Mean Reporting rate: SABAP2	Preferred Habitat	Potential Likelihood of Occurrence
<i>Gyps coprotheres</i> (Cape Vulture)	Endangered	Endangered	17.16	46.0 (for pentad 2600_2605)	Mainly confined to mountain ranges, especially near breeding site. Ventures far afield in search of food.	A regular foraging/scavenging visitor to the study site pending the presence of food (e.g. livestock carcasses).
<i>Gyps africanus</i> (White-backed Vulture)	Critically Endangered	Critically Endangered	16.18	46.7 (for pentad 2600_2605)	Breed on tall, flat-topped trees. Mainly restricted to large rural or game farming areas.	A regular foraging/scavenging visitor to the study site pending the presence of food (e.g. livestock carcasses).
<i>Leptoptilos crumeniferus</i> (Marabou Stork)	-	Near threatened	0.70	1.56	Varied, from savanna to wetlands, pans and floodplains – dependant of game farming areas	An irregular scavenging visitor to the area.
<i>Polemaetus bellicosus</i> (Martial Eagle)	Endangered	Endangered	-	4.5 (for pentad 2605_2605)	Varied, from open karroid shrub to lowland savanna.	An irregular foraging visitor. It was last recorded from pentad 2605_2605 south-east of the study site on 28 Jan 2012.
<i>Sagittarius serpentarius</i> (Secretarybird)	Endangered	Vulnerable	2.45	1.56 (for pentad 2605_2605)	Prefers open grassland or lightly wooded habitat.	Regarded as an irregular foraging visitor to the study site despite the widespread presence of

Species	Global Conservation Status*	National Conservation Status**	Mean Reporting rate: SABAP1	Mean Reporting rate: SABAP2	Preferred Habitat	Potential Likelihood of Occurrence
						suitable foraging habitat.
<i>Torgos tracheliotos</i> (Lapped-faced Vulture)	Endangered	Endangered	5.63	26.7 (for pentad 2600_2605)	Lowveld and Kalahari savanna; mainly on game farms and reserves	A regular foraging/scavenging visitor to the study site pending the presence of food (e.g. livestock carcasses).

3.8 Preliminary avifaunal sensitivity

A preliminary sensitivity map was compiled, illustrating habitat units comprising of potential sensitive elements based on the following arguments (Figure 7):

Areas of medium sensitivity

It includes the artificial livestock watering points, moist grassland and extensive open grassland and bush clump mosaics. The bush clumps, especially the taller *Vachellia erioloba* is often used as roosting platforms for vultures.

The artificial livestock watering points attracted large numbers of granivore passerine and non-passerine bird species, of which many need to drink water on a daily basis. The placement of electrical infrastructure in close proximity to these areas could increase potential avian collisions with the infrastructure. These areas are therefore of artificial origin, but could be relocated to other areas.

The extensive open grassland and bush clump mosaics provide potential suitable foraging habitat for some collision-prone bird species, including the Northern Black Korhaan (*Afrotis afroides*) with the potential to interact (e.g. collide) with the proposed electrical infrastructure. However, reporting rates for threatened and near threatened bird species are anticipated to be relatively low, thereby suggesting a medium sensitivity rating instead of a high sensitivity even though the majority of the habitat is natural. In addition, the open grassland and bush clump mosaics are widespread in the region.

Areas of low sensitivity

These habitat units are represented by transformed types and include a small excavated area.

The preliminary sensitivity map shows a large surface area that is earmarked with medium sensitivity. There is a probability that some of these units or part thereof could

have higher (or lower) sensitivity ratings. It is therefore expected that some of the units or part thereof could represent different sensitivity ratings to those displayed in Figure 7 pending the outcome of a detailed season surveys.




Figure 7: A map illustrating the preliminary avifaunal sensitivity of the area based on habitat types supporting bird taxa of conservation concern and important ecological function.

3.9 Overview of Avian Impacts at Solar Facilities

3.9.1 Background to solar facilities and their impact on birds

Birds are mobile, and are therefore also more readily affected by solar facilities than other taxonomic groups (e.g. mobile mammals that could move away from the facilities due to displacement). In fact, birds are also vulnerable to impacts caused by other types of energy facilities such as overhead power lines and wind farms. Little information is available on the impacts of solar energy facilities on birds although Gunerhan *et al.* (2009), McCrary *et al.* (1986), Tsoutsos *et al.* (2005) and the recent investigation reports on bird fatalities in the USA by Kagen *et al.* (2014) and Walston *et al.* (2016) provide discussions thereof. These studies have shown that avian fatalities vary greatly between the geographic positions of the solar facilities and also depend on the type of solar facility. In addition, very few of the large solar facilities in operation undertake systematic monitoring of avian fatalities, which explains the lack of detailed information of avian impacts. According to these studies conducted at both

Concentrated Solar Power (CSP) and PV facilities, avian incidental fatalities range from 14 to over 180 birds which were summarised over a survey period conducted during one to three years. According to the Walston *et al.* (2016) assessment, the average annual mortality rate for known utility-scale solar facilities (the annual number of estimated bird deaths per megawatt of electrical capacity) is 2.7, and 9.9 for known and unknown fatalities (which include carcasses found on the project site of which the death is not known). McCrary *et al.* (1986) found an average rate of mortality of 1.9-2.2 birds per week affecting 0.6-0.7 % of the local bird population. However, most of the avian fatalities at these solar facilities are also probably underestimated since 10-30 % of dead birds are removed by scavengers before being noted.. From these analyses and assessments it was evident that:

- Medium levels of bird fatalities occur at PV sites when compared to CSP sites (when taking powerline collisions into account).
- Approximately 81 % of all avian mortalities were caused by collisions, including collisions with electrical distribution lines.
- Most of the mortalities were small passerines (especially swallows).
- Fatalities at these solar facilities also include waterbirds (e.g. grebes, herons and gulls) which were probably attracted by the apparent "lake effect" caused by the reflective surface of the PV panels.
- Approximately 10-11 % of the fatalities consists of waterbirds, but could be as high as 49 % at certain facilities.
- It is unclear if the "lake effect" caused by the panels (at PV facilities) or mirrors (at CSP facilities) are the main cause of birds colliding or interacting with the infrastructure (since both waterbirds and other passerines are colliding with the infrastructure).
- Most of the fatalities are of resident birds as opposed to migratory species.

In a review report by Harrison *et al.* (2016), an attempt was made to provide evidence of the impacts caused by solar PV facilities alone (not combined with CSP facilities) on birds in the UK. These authors reviewed approximately 420 scientific documents, including 37 so-called "grey" literature from non-government and government organisations for any evidence relating to the ecological impacts of solar PV facilities. Their main findings were as follows:

- The majority of the documents were not relevant and peer-reviewed documents of experimental scientific evidence on avian fatalities were non-existent.
- Results based on carcass searches suggest that the bird collision risk at PV developments are low, although these studies did not take collision by overhead power lines into account.
- Many of the documents recommended that PV developments in close proximity to protected areas should be avoided.
- The PV panels reflect polarised light, which can attract polarotactic insects with potential impact to their reproductive biology. In addition, the polarising effect

of the PV panels may also induce drinking behaviour in some birds, which may mistake the panels for water.

They conclude that impact assessment reports should consider taxon-specific requirements of birds and their guilds.

3.9.2 Potential impacts of PV solar facilities on birds

The magnitude and significance of impacts to birds caused by solar facilities will depend on the following factors:

- The geographic locality of the planned solar facility;
- The size or surface extent of the solar facility;
- The type of solar facility (according to the technologies applied, e.g. PV or Concentrated Solar Power (CSP)); and
- The occurrence of collision-prone bird species (which are often closely related to the locality of the solar facility).

Any planned solar facility corresponding to an area with many threatened, range-restricted or collision-prone species will have a higher impact on these birds. In addition, any planned solar facility located in close proximity to important flyways, wetland systems or roosting/nesting sites used by the aforementioned species will have a higher impact.

The main impacts associated with PV solar facilities include (Jenkins *et al.*, 2017):

- The loss of habitat and subsequent displacement of bird species due to the ecological footprint required during construction;
- Disturbances caused to birds during construction and operation;
- Direct interaction (collision trauma) by birds with the surface infrastructure (photovoltaic panels) caused by polarised light pollution and/or waterbirds colliding with the panels (as they are mistaken for waterbodies);
- Collision with associated infrastructure (mainly overhead powerlines and reticulation);
- Attracting novel species to the area (owing to the artificial provision of new habitat such as perches and shade) which could compete with the residing bird population.

3.10 Potential Impacts associated with the Euphorbia PV, Hillardia PV and Verbena PV Solar Energy Facilities

3.10.1 Loss of habitat and displacement of birds

Most of the study site will be cleared of vegetation and habitat to accommodate the panel arrays and associated infrastructure. Clearing of vegetation will inevitably result in the loss of habitat and displacement of bird species. From the preliminary results it is evident that large-bodied species are more likely to become displaced as opposed to small passerine species. It is particularly biome-restricted, endemic and conservation

important species that are likely to become displaced, as well as habitat specialists (e.g. grassland specialists) which will disappear from the area. These include mainly passerine and smaller non-passerine species inhabiting the untransformed dolomite grasslands and bush clump mosaics.

To quantify the impact it is necessary to calculate the number of birds (density) lost or displaced by the activity, including estimated density values of important species per unit area of habitat. This will be conducted during an austral summer season survey of the proposed PV facilities. From a preliminary analysis, the following bird species are most likely to be impacted by the loss of habitat due to their habitat requirements, fecundity and conservation status (although not limited to) due to the proposed development:

- Northern Black Korhaan (*Afrotis afraoides*);
- Ashy Tit (*Melaniparus cinerascens*);
- Kalahari Scrub Robin (*Cercotrichas paena*);
- Orange River Francolin (*Scleroptila gutturalis*) and potentially also small to medium birds of prey such as:
 - Black-winged Kite (*Elanus caeruleus*);
 - Gabar Goshawk (*Micronisus capensis*);
 - Amur Falcon (*Falco amurensis*);
 - Lesser Kestrel (*Falco rupicolus*) and
 - Black-chested Snake-eagle (*Circaetus pectoralis*).

4.10.2 Collision trauma caused by photovoltaic panels (the "lake-effect")

The study site is not located in close proximity to any major wetland system or water body. The nearest large wetland system that is inundated is approximately 9 km from the study site, which explain the low expected occurrence of waterbird taxa at the study site. The waterbirds could accidentally mistake the reflective panels for waterbodies, thereby resulting in bird collisions with the panel surfaces. At this stage the impact is considered to be low, depending on subsequent site visits (e.g. pre-construction monitoring).

However, desktop results and previous site observations show that the following species could potentially interact with the panel infrastructure:

- Yellow-billed Duck (*Anas undulata*);
- Red-billed Teal (*Anas erythrorhynchus*);
- South African Shelduck (*Tadorna cana*);
- Spur-winged Goose (*Plectropterus gambiensis*);
- Egyptian Goose (*Alopochen aegyptiaca*);
- Black-headed Heron (*Ardea melanocephala*); and probably also
- Grey Heron (*Ardea cinerea*) and
- White-faced Duck (*Dendrocygna viduata*).

Of these species, the Spur-winged Goose, Egyptian Goose and Black-headed Heron were regarded as the only regular visitor to the immediate surroundings.

4.10.3 Creation of "new" avian habitat and bird pollution

It is possible that the infrastructure (during operation) could attract bird species which may occupy the site or interact with the local bird assemblages in the wider region. These include alien and cosmopolitan species, as well as aggressive omnivorous passerines which could displace other bird species from the area:

- House Sparrow (*Passer domesticus*);
- Common Myna (*Acridotheres tristis*);
- Pied Crow (*Corvus albus*); and
- Speckled Pigeon (*Columba guinea*).

The infrastructure may attract large numbers of roosting columbid taxa, especially Speckled Pigeons (*Columba guinea*), which may result in avian "pollution" through excreta, thereby fouling the panel surfaces. The impact is manageable.

3.10.4 Interaction with overhead powerlines (grid connection)

An overhead powerline is proposed for the grid connection from the facility substation to the Watershed Main Transmission Substation. Birds are impacted in three ways by means of overhead powerlines (described below). It is however a common rule that large and heavy-bodied terrestrial bird species are more at risk of being affected in a negative way when interacting with powerlines in general. These include the following:

- *Electrocution*

Electrocution happens when a bird bridges the gap between the live components or a combination of a live and earth component of a power line, thereby creating a short circuit. This happens when a bird, mainly a species with a fairly large wingspan attempts to perch on a tower or attempts to fly-off a tower. Many of these species include vultures (of the genera *Gyps* and *Torgos*) as well as other large birds of prey such as the Martial Eagle (*Polemaetus bellicosus*) (Ledger & Annegarn, 1981; Kruger, 1999; Van Rooyen, 2000). These species will attempt to roost and even breed on the tower structures if available nesting platforms are a scarce commodity in the area. Other types of electrocutions happen by means of so-called "bird-streamers". This happens when a bird, especially when taking off, excretes and thereby causes a short-circuit through the fluidity excreta (Van Rooyen & Taylor, 1999).

Large transmission lines (from 220 kV to 765 kV) are seldom a risk of electrocution, although smaller distribution lines (88 – 132kV) pose a higher risk. However, for this project, the design of the pylon is an important consideration in preventing bird

electrocutions. **The proposed pylon design must incorporate the following design parameters:**

- The clearances between the live components should exceed the wingspan of any bird species;
- The height of the tower should allow for unrestricted movement of terrestrial birds between successive pylons;
- The live components should be “bundled” to increase the visibility for approaching birds;
- “Bird streamers” should be eliminated by discouraging birds from perching above the conductors.

It is therefore recommended that the pylon design incorporates “features as illustrated by Figure 8⁶.”

From Figure 8 it is clear that perching of birds is discouraged by the addition of diagonal crossbars or by doing away with the crossbars that holds the conductors in place. Bird “streamers” are also eliminated by fitting the poles with bird guards/spikes above the conductors. However, safe perching is facilitated by the fitment of a horizontal bar on top of the pole structure without the risk of electrocution (due to the perpendicular orientation of the bar relative to the conductors).




Figure 8: Two bird-friendly tower designs to be used for the current project.

⁶ Please note that these are examples of recommended pylon designs. These are taken from steel monopole pylons.

- *Collision*

Collisions with earth wires have probably accounted for most bird-powerline interactions in South Africa. In general, the earth wires are much thinner in diameter when compared to the live components, and therefore less visible to approaching birds. Many of the species likely to be affected include heavy, large-bodied terrestrial species such as bustards, korhaans and a variety of waterbirds that are not very agile or manoeuvrable once airborne. These species, especially those with the habit of flying with outstretched necks (e.g. most species of storks) find it difficult to make a sudden change in direction while flying – resulting in the bird flying into the earth wires.

Areas where bird collisions are likely to be high could be ameliorated by marking the lines with appropriate bird deterrent devices such as “bird diverters” and “flappers” to increase the visibility of the lines. Table 4 provides a list of potential bird species that could collide with the overhead grid connection

- *Physical disturbances and habitat destruction caused during construction and maintenance*

It is anticipated that part of the power line servitude will be cleared of vegetation. In addition, construction activities go hand in hand with high ambient noise levels. Although construction is considered temporary, many species will vacate the area during the construction phase and will become temporarily displaced.

The artificial livestock watering points also deserve special consideration since these features are often overlooked or neglected during the construction of power lines as they often attract large numbers of small passerine birds and birds of prey (the latter often include falconiform taxa which hunt small passerines). Construction activities in close proximity to these features could possibly displace these individuals from the area or increase the risk of collision. Nevertheless, these features could easily be removed or relocated to other areas.

It is important to note that impacts associated with the proposed powerline infrastructure will be assessed as part of a standalone Basic Assessment process and not as part of the facility EIAs. This baseline report does however report on the potential impacts as the two (the facilities and the grid connection infrastructure) are co-dependant and should be viewed holistically.

3.11 Collision-prone bird species

A total of 41 collision-prone bird species have been recorded in the wider study area (*sensu* SABAP2), of which nine species could interact with the PV panels (Table 4). However, most of these species are more at risk of colliding with overhead powerlines (to be assessed during a separate Basic Assessment process).

Table 4: Collision-prone bird species and Red listed species (in red) expected to be present on the study site.

Common Name	Scientific Name	National conservation status (sensu Taylor et al., 2015)
Abdim's Stork	<i>Ciconia abdimii</i>	NT
African Hawk-eagle	<i>Aquila spilogaster</i>	
African Harrier-Hawk	<i>Polyboroides typus</i>	
African Sacred Ibis	<i>Threskiornis aethiopicus</i>	
Amur Falcon	<i>Falco amurensis</i>	
Black Kite	<i>Milvus migrans</i>	
Black-chested Snake Eagle	<i>Circaetus pectoralis</i>	
Black-headed Heron	<i>Ardea melanocephala</i>	
Black-winged Kite	<i>Elanus caeruleus</i>	
Brown Snake Eagle	<i>Circaetus cinereus</i>	
Cape Vulture	<i>Gyps coprotheres</i>	EN
Common (Steppe) Buzzard	<i>Buteo buteo vulpinus</i>	
Coqui Francolin	<i>Peliperdix coqui</i>	
Egyptian Goose	<i>Alopochen aegyptiaca</i>	
Gabar Goshawk	<i>Micronisus capensis</i>	
Greater Kestrel	<i>Falco rupicoloides</i>	
Hadada Ibis	<i>Bostrychia hagedash</i>	
Helmeted Guineafowl	<i>Numida meleagris</i>	
Lanner Falcon	<i>Falco biarmicus</i>	VU
Lappet-faced Vulture	<i>Torgos tracheliotos</i>	EN
Lesser Kestrel	<i>Falco naumanni</i>	
Marsh Owl	<i>Asio capensis</i>	
Martial Eagle	<i>Polemaetus bellicosus</i>	EN
Northern Black Korhaan	<i>Afrotis afrooides</i>	
Orange River Francolin	<i>Scleroptila gutturalis</i>	
Pale Chanting Goshawk	<i>Melierax canorus</i>	
Pied Crow	<i>Corvus albus</i>	
Red-footed Falcon	<i>Falco vespertinus</i>	NT
Rock Dove	<i>Columba livia</i>	
Rock Kestrel	<i>Falco rupicolus</i>	
Secretarybird	<i>Sagittarius serpentarius</i>	VU
Speckled Pigeon	<i>Columba guinea</i>	
Spotted Eagle-Owl	<i>Bubo africanus</i>	
Spur-winged Goose	<i>Plectropterus gambensis</i>	
Swainson's Spurfowl	<i>Pternistis swainsonii</i>	
Western Barn Owl	<i>Tyto alba</i>	
Western Cattle Egret	<i>Bubulcus ibis</i>	
White-backed Vulture	<i>Gyps africanus</i>	CR
Yellow-billed Duck	<i>Anas undulata</i>	
Yellow-billed Kite	<i>Milvus aegyptius</i>	

The study site does not coincide with any prominent wetland system or impoundment which will lower the risk of waterbird collisions with the proposed electrical infrastructure.

3.11.1 Vultures

Three species of vulture occur in the study area, which are prone towards electrocution and collision with powerlines⁷. These include the globally critically endangered White-backed Vulture (*Gyps africanus*), the globally endangered Cape Vulture (*G. coprotheres*) and the globally endangered Lapped-faced Vulture (*Torgos tracheliotos*). These species are of international significance and any mortality of adult individuals could have a negative effect on its species' population recruitment. Most of these suffer from a shortage of food supplies which is responsible for low reproductive rates, especially for Cape Vultures (Taylor *et al.*, 2015). In addition, most of these species also tend to congregate at mammalian carcasses, where they feed in large groups, especially in terms of Cape Vultures. In addition, Cape Vultures also typically search for food in groups. It is such congregations which increase the risk of mortalities whenever these individuals forage or roost in close proximity to powerlines. For example, the proposed study area coincides with the foraging rangeland of Cape Vultures as evidenced by dispersal data obtained from vulture individuals fitted with satellite tracking devices (Figure 9).

The highest number of mortalities due to electrocution and collision recorded in the study region pertains to Cape Vultures (*Gyps coprotheres*) and White-backed Vultures (*Gyps africanus*) (according to the electrical infrastructure mortality incident register) (Figure 10). Most of the mortalities were caused during electrocution from smaller distribution lines in the area, although a significant number of Cape Vulture mortalities (c. 30 %) were also caused by collisions with transmission lines (Figure 11). There is a definite correlation between the size (in terms of voltage) of the powerline and the type of mortality, whereby electrocution incidents were prominent from distribution lines, while collisions were caused by transmission lines. Most of the powerline interactions also occurred in the Ventersdorp and Lichtenburg area (Figure 12), with a single mass mortality involving 10 Cape Vultures and eight White-backed Vultures on 09 March 2009. **It clearly shows that when these species congregate (for example when feeding from a carcass in close proximity to an overhead powerline or when roosting on pylons or nearby structures in close proximity to powerlines), the risk of mortality due to both electrocution and collision is greatly increased.**

⁷ It is important to note that impacts associated with the proposed powerline infrastructure will be assessed as part of a standalone Basic Assessment process and not as part of the facility EIAs. This baseline report does however report on powerline collision prone species as the two (the facilities and the grid connection infrastructure) are co-dependant and should be viewed holistically.




Figure 9: The occurrence of Cape Vultures (*Gyps coprotheres*) within the study region that were fitted with satellite trackers.




Figure 10: The number of mortalities (electrocutions and collisions) per bird species due to transmission and distribution lines in the study area (1996-2018).




Figure 11: The number of mortalities per bird species caused by electrocutions (distribution lines) and collisions (transmission lines) (1996-2018).




Figure 12: The number of bird mortalities caused by power lines per geographic locality (1996-2018), including the Lichtenburg area.

4. PLAN OF STUDY FOR THE EIA PHASE

Due to the limited level of detail that is normally implemented during a scoping assessment, it is imperative that detailed avifaunal investigations be conducted on the study area which includes seasonal coverage.

4.1 Proposed approach and methods

The following methods are proposed during an austral summer season survey:

- Active searching and the compilation of a bird inventory while traversing much of the available habitat types;
- The determination of the occurrence of Red Data species and collision-prone bird species;
- The identification and mapping of suitable habitat for species of conservation concern while focussing on structural and topographical cues;
- A landscape analysis of important flyways or daily flight paths corresponding to important landscape features; and
- Density estimates will be collected by means of point counts to evaluate the dominant/typical species and their respective relative densities at each site. At each point the number of bird species seen will be recorded, as well as their respective abundances and distance from the observer (by means of a rangefinder). The data generated from the point counts will be analysed according to Clarke & Warwick (1994) based on the computed percentage contribution (%) of each species including the consistency (calculated as the similarity coefficient/standard deviation) of its contribution to the each habitat type.

5. REFERENCES

Birdlife South Africa. 2018. *BirdLife South Africa Checklist of Birds in South Africa*, 2018.

Clarke, K.R. & Warwick, R.M. 1994. *Changes in marine communities: An approach to statistical analysis and interpretation*. Natural Environmental Research Council, United Kingdom.

Convention on Biological Diversity. Signed 1993 and ratified 2 November 1995.

Geoterrainimage. 2015. *The South African National Land cover Dataset*. Version 05.

Gill, F, D Donsker, & P Rasmussen (Eds). 2021. IOC World Bird List (v 11.2). Doi 10.14344/IOC.ML.10.2. <http://www.worldbirdnames.org/>.

Gunerhan, H., Hepbasli, A. & Giresunlu, U. 2009. Environmental impacts from the solar energy systems. *Energy Sources, Part A: Recovery, Utilization and Environmental Effects* 31: 131-138.

Hardaker, T. 2018. Southern African Bird List - Version 08 - 11 March 2018.

Harrison, C., Lloyd, H. & Field, C. 2016. *Evidence review of the impact of solar farms on birds, bats and general ecology*. NEER012 report, Manchester Metropolitan University, UK.

Harrison, J.A., Allan, D.G., Underhill, L.G., Herremans, M., Tree, A.J., Parker, V. & Brown, C.J. (eds.). 1997. *The Atlas of Southern African Birds. Vol. 1 & 2*. BirdLife South Africa, Johannesburg.

Hockey, P.A.R., Dean, W.R.J. & Ryan, P.G. (eds.) 2005. *Roberts – Birds of Southern Africa*, VIIth ed. The Trustees of the John Voelker Bird Book Fund, Cape Town.

IUCN Red List of Threatened Species. Version 2021. <http://www.iucnredlist.org/>.

Jenkins, A.R, Ralston-Paton, S & Smit-Robinson, H.A. 2017. Best practice guidelines: Birds and Solar Energy. Guidelines for assessing and monitoring the impact of solar power generating facilities on birds in southern Africa. BirdLife South Africa.

Kagen, R.A., Verner, T.C., Trail, PW & Espinoza, E.O. 2014. Avian mortality at solar energy facilities in southern California: A preliminary analysis. Unpublished report by the National Fish and Wildlife Forensics Laboratory, USA.

Kruger, R. 1999. *Towards solving raptor electrocutions on Eskom Distribution Structures in South Africa*. M. Phil. Mini-thesis. University of the Orange Free State. Bloemfontein. South Africa.

Ledger, J. & Annegarn, H.J. 1981. Electrocution Hazards to the Cape Vulture (*Gyps coprotheres*) in South Africa. *Biological Conservation* 20: 15-24.

Marnewick, M.D., Retief, E.F., Theron, N.T., Wright, D.R. And Anderson, T.A. 2015. *Important Bird and Biodiversity Areas of South Africa*. Johannesburg: BirdLife South Africa.

McCrary, M.D., McKernan, R.L., Schreiber, R.W., Wagner, W.D. & Sciarotta, T.C. 1986. Avian mortality at a solar energy power plant. *Journal of Field Ornithology* 57: 135-141.

Mucina, L. & Rutherford, M.C. (eds.). 2006. The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia* 19. South African National Biodiversity Institute, Pretoria.

National Environmental Management Biodiversity Act, 2004 (Act No. 10 of 2004).

Pachnoda Consulting. 2018. Development of the Lichtenburg 2 PV solar energy facility and associated infrastructure on a site near Lichtenburg, North West Province. A report compiled for Savanna Environmental.

Pachnoda Consulting. 2012. Development of the Barleria PV solar energy facility and associated infrastructure on a site near Lichtenburg, North West Province. A report compiled for Savanna Environmental.

Taylor, M.R., Peacock, F. & Wanless, R. (eds.). 2015. *The Eskom Red Data Book of Birds of South Africa, Lesotho and Swaziland*. BirdLife South Africa, Johannesburg

Tsoutsos, T., Frantzeskaki, N. & Gekas, V. 2005. Environmental impacts from solar energy technologies. *Energy Policy* 33: 289-296.

Van Rooyen, C.S. 2000. An overview of Vulture Electrocutions in South Africa. *Vulture News* 43: 5-22.

Van Rooyen, C.S. & Taylor, P.V. 1999. *Bird streamers as probable cause of electrocutions in South Africa*. EPRI Workshop on Avian Interactions with Utility Structures, Charleston, South Carolina.

Vosloo, H. 2003. Birds and power lines. *ESI Africa* 3: 38.

Walston Jr. L.J., Rollins, K.E., LaGory, K.E., Smith, K.P. & Meyers, S.A. 2016. A preliminary assessment of avian mortality at utility-scale solar energy facilities in the United States. *Renewable Energy* 92 (2016) 405-414.

www.sabap2.birdmap.africa

Appendix 1: A shortlist of bird species expected to be present on the study area. The list provides an indication of the species occurrence according to SABAP2 reporting rates.

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
78	Abdim's Stork	<i>Ciconia abdimii</i>	0.00	0	3.70	1
432	Acacia Pied Barbet	<i>Tricholaema leucomelas</i>	37.33	28	3.70	1
171	African Harrier-Hawk	<i>Polyboroides typus</i>	2.67	2	0.00	0
141	African Hawk-eagle	<i>Aquila spilogaster</i>	-	1	-	-
418	African Hoopoe	<i>Upupa africana</i>	44.00	33	0.00	0
387	African Palm Swift	<i>Cypsiurus parvus</i>	38.67	29	0.00	0
682	African Paradise Flycatcher	<i>Terpsiphone viridis</i>	8.00	6	0.00	0
692	African Pipit	<i>Anthus cinnamomeus</i>	37.33	28	3.70	1
544	African Red-eyed Bulbul	<i>Pycnonotus nigricans</i>	42.67	32	0.00	0
606	African Reed Warbler	<i>Acrocephalus baeticatus</i>	20.00	15	0.00	0
81	African Sacred Ibis	<i>Threskiornis aethiopicus</i>	10.67	8	0.00	0
576	African Stonechat	<i>Saxicola torquatus</i>	44.00	33	0.00	0
247	African Wattled Lapwing	<i>Vanellus senegallus</i>	1.33	1	0.00	0
772	Amethyst Sunbird	<i>Chalcomitra amethystina</i>	5.33	4	0.00	0
119	Amur Falcon	<i>Falco amurensis</i>	21.33	16	3.70	1
575	Ant-eating Chat	<i>Myrmecocichla formicivora</i>	46.67	35	7.41	2
533	Arrow-marked Babbler	<i>Turdoides jardineii</i>	2.67	2	0.00	0
514	Ashy Tit	<i>Melaniparus cinerascens</i>	4.00	3	0.00	0
510	Banded Martin	<i>Riparia cincta</i>	13.33	10	3.70	1
493	Barn Swallow	<i>Hirundo rustica</i>	36.00	27	7.41	2
622	Bar-throated Apalis	<i>Apalis thoracica</i>	1.33	1	0.00	0
128	Black Kite	<i>Milvus migrans</i>	1.33	1	0.00	0

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
712	Black-backed Puffback	<i>Dryoscopus cubla</i>	1.33	1	0.00	0
650	Black-chested Prinia	<i>Prinia flavicans</i>	68.00	51	7.41	2
146	Black-chested Snake Eagle	<i>Circaetus pectoralis</i>	4.00	3	0.00	0
431	Black-collared Barbet	<i>Lybius torquatus</i>	34.67	26	3.70	1
715	Black-crowned Tchagra	<i>Tchagra senegalus</i>	2.67	2	0.00	0
55	Black-headed Heron	<i>Ardea melanocephala</i>	26.67	20	3.70	1
521	Black-headed Oriole	<i>Oriolus larvatus</i>	2.67	2	0.00	0
245	Blacksmith Lapwing	<i>Vanellus armatus</i>	70.67	53	3.70	1
860	Black-throated Canary	<i>Crithagra atrogularis</i>	42.67	32	0.00	0
130	Black-winged Kite	<i>Elanus caeruleus</i>	32.00	24	25.93	7
839	Blue Waxbill	<i>Uraeginthus angolensis</i>	25.33	19	3.70	1
722	Bokmakierie	<i>Telophorus zeylonus</i>	44.00	33	3.70	1
145	Brown Snake Eagle	<i>Circaetus cinereus</i>	1.33	1	0.00	0
714	Brown-crowned Tchagra	<i>Tchagra australis</i>	13.33	10	7.41	2
402	Brown-hooded Kingfisher	<i>Halcyon albiventris</i>	1.33	1	0.00	0
731	Brubru	<i>Nilaus afer</i>	4.00	3	3.70	1
695	Buffy Pipit	<i>Anthus vaalensis</i>	1.33	1	0.00	0
4131	Burchell's Coucal	<i>Centropus burchellii</i>	17.33	13	0.00	0
703	Cape Longclaw	<i>Macronyx capensis</i>	30.67	23	3.70	1
531	Cape Penduline Tit	<i>Anthoscopus minutus</i>	2.67	2	0.00	0
581	Cape Robin-Chat	<i>Cossypha caffra</i>	21.33	16	0.00	0
786	Cape Sparrow	<i>Passer melanurus</i>	74.67	56	14.81	4
737	Cape Starling	<i>Lamprotornis nitens</i>	29.33	22	7.41	2
316	Ring-necked Dove	<i>Streptopelia capicola</i>	24.00	18	14.81	4
106	Cape Vulture	<i>Gyps coprotheres</i>	10.67	8	0.00	0

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
686	Cape Wagtail	<i>Motacilla capensis</i>	57.33	43	0.00	0
799	Cape Weaver	<i>Ploceus capensis</i>	5.33	4	0.00	0
1172	Cape White-eye	<i>Zosterops virens</i>	28.00	21	0.00	0
568	Capped Wheatear	<i>Oenanthe pileata</i>	9.33	7	0.00	0
484	Chestnut-backed Sparrow-Lark	<i>Eremopterix leucotis</i>	10.67	8	11.11	3
658	Chestnut-vented Warbler	<i>Curruca subcoerulea</i>	38.67	29	7.41	2
673	Chinspot Batis	<i>Batis molitor</i>	6.67	5	0.00	0
872	Cinnamon-breasted Bunting	<i>Emberiza tahapisi</i>	9.33	7	3.70	1
631	Cloud Cisticola	<i>Cisticola textrix</i>	17.33	13	3.70	1
196	Common Buttonquail	<i>Turnix sylvaticus</i>	-	1	-	-
154	Common (Steppe) Buzzard	<i>Buteo buteo vulpinus</i>	4.00	3	7.41	2
734	Common Myna	<i>Acridotheres tristis</i>	69.33	52	7.41	2
189	Common Quail	<i>Coturnix coturnix</i>	n/a			
421	Common Scimitarbill	<i>Rhinopomastus cyanomelas</i>	16.00	12	0.00	0
843	Common Waxbill	<i>Estrilda astrild</i>	18.67	14	0.00	0
594	Common Whitethroat	<i>Curruca communis</i>	1.33	1	0.00	0
173	Coqui Francolin	<i>Peliperdix coqui</i>	6.67	5	0.00	0
439	Crested Barbet	<i>Trachyphonus vaillantii</i>	65.33	49	0.00	0
711	Crimson-breasted Shrike	<i>Laniarius atrococcineus</i>	17.33	13	0.00	0
242	Crowned Lapwing	<i>Vanellus coronatus</i>	70.67	53	7.41	2
545	Dark-capped Bulbul	<i>Pycnonotus tricolor</i>	32.00	24	0.00	0
630	Desert Cisticola	<i>Cisticola aridulus</i>	20.00	15	7.41	2
352	Diederik Cuckoo	<i>Chrysococcyx caprius</i>	32.00	24	3.70	1
1183	Eastern Clapper Lark	<i>Mirafrja fasciolata</i>	18.67	14	0.00	0
89	Egyptian Goose	<i>Alopochen aegyptiaca</i>	20.00	15	0.00	0

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
404	European Bee-eater	<i>Merops apiaster</i>	28.00	21	0.00	0
678	Fairy Flycatcher	<i>Stenostira scita</i>	n/a			
570	Familiar Chat	<i>Oenanthe familiaris</i>	4.00	3	0.00	0
665	Fiscal Flycatcher	<i>Melaenornis silens</i>	42.67	32	7.41	2
517	Fork-tailed Drongo	<i>Dicrurus adsimilis</i>	1.33	1	3.70	1
162	Gabar Goshawk	<i>Micronisus capensis</i>	-	1	-	-
874	Golden-breasted Bunting	<i>Emberiza flaviventris</i>	2.67	2	3.70	1
447	Golden-tailed Woodpecker	<i>Campethera abingoni</i>	2.67	2	0.00	0
785	Great Sparrow	<i>Passer motitensis</i>	2.67	2	0.00	0
440	Greater Honeyguide	<i>Indicator indicator</i>	4.00	3	0.00	0
122	Greater Kestrel	<i>Falco rupicoloides</i>	4.00	3	3.70	1
502	Greater Striped Swallow	<i>Cecropis cucullata</i>	48.00	36	0.00	0
419	Green Wood Hoopoe	<i>Phoeniculus purpureus</i>	9.33	7	0.00	0
830	Green-winged Pytilia	<i>Pytilia melba</i>	10.67	8	3.70	1
339	Grey Go-away-bird	<i>Crinifer concolor</i>	20.00	15	0.00	0
557	Groundscraper Thrush	<i>Turdus litsitsirupa</i>	6.67	5	0.00	0
84	Hadada Ibis	<i>Bostrychia hagedash</i>	62.67	47	0.00	0
192	Helmeted Guineafowl	<i>Numida meleagris</i>	44.00	33	11.11	3
784	House Sparrow	<i>Passer domesticus</i>	54.67	41	11.11	3
835	Jameson's Firefinch	<i>Lagonosticta rhodopareia</i>	2.67	2	0.00	0
586	Kalahari Scrub Robin	<i>Cercotrichas paena</i>	34.67	26	7.41	2
1104	Karoo Thrush	<i>Turdus smithi</i>	54.67	41	0.00	0
552	Kurrichane Thrush	<i>Turdus libonyana</i>	n/a			
114	Lanner Falcon	<i>Falco biarmicus</i>	4.00	3	0.00	0
108	Lappet-faced Vulture	<i>Torgos tracheliotos</i>	5.33	4	0.00	0

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
317	Laughing Dove	<i>Spilopelia senegalensis</i>	90.67	68	25.93	7
706	Lesser Grey Shrike	<i>Lanius minor</i>	14.67	11	0.00	0
442	Lesser Honeyguide	<i>Indicator minor</i>	4.00	3	0.00	0
125	Lesser Kestrel	<i>Falco naumanni</i>	16.00	12	3.70	1
413	Lilac-breasted Roller	<i>Coracias caudatus</i>	1.33	1	0.00	0
410	Little Bee-eater	<i>Merops pusillus</i>	8.00	6	3.70	1
385	Little Swift	<i>Apus affinis</i>	34.67	26	3.70	1
621	Long-billed Crombec	<i>Sylvietta rufescens</i>	9.33	7	0.00	0
852	Long-tailed Paradise Whydah	<i>Vidua paradisaea</i>	5.33	4	0.00	0
818	Long-tailed Widowbird	<i>Euplectes progne</i>	41.33	31	14.81	4
661	Marico Flycatcher	<i>Melaenornis mariquensis</i>	5.33	4	0.00	0
381	Marsh Owl	<i>Asio capensis</i>	-	1	-	-
607	Marsh Warbler	<i>Acrocephalus palustris</i>	5.33	4	3.70	1
142	Martial Eagle	<i>Polemaetus bellicosus</i>	1.33	1	0.00	0
456	Melodious Lark	<i>Mirafra cheniana</i>	n/a			
564	Mountain Wheatear	<i>Myrmecocichla monticola</i>	2.67	2	0.00	0
318	Namaqua Dove	<i>Oena capensis</i>	16.00	12	0.00	0
637	Neddicky	<i>Cisticola fulvicapilla</i>	18.67	14	3.70	1
1035	Northern Black Korhaan	<i>Afrotis afraoides</i>	34.67	26	3.70	1
179	Orange River Francolin	<i>Scleroptila gutturalis</i>	20.00	15	3.70	1
838	Orange-breasted Waxbill	<i>Amandava subflava</i>	4.00	3	0.00	0
165	Pale Chanting Goshawk	<i>Melierax canorus</i>	n/a			
522	Pied Crow	<i>Corvus albus</i>	45.33	34	7.41	2
746	Pied Starling	<i>Lamprotornis bicolor</i>	6.67	5	7.41	2
846	Pin-tailed Whydah	<i>Vidua macroura</i>	26.67	20	0.00	0

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
694	Plain-backed Pipit	<i>Anthus leucophrys</i>	4.00	3	0.00	0
844	Quailfinch	<i>Ortygospiza atricollis</i>	18.67	14	3.70	1
642	Rattling Cisticola	<i>Cisticola chiniana</i>	10.67	8	0.00	0
708	Red-backed Shrike	<i>Lanius collurio</i>	22.67	17	0.00	0
837	Red-billed Firefinch	<i>Lagonosticta senegala</i>	14.67	11	0.00	0
805	Red-billed Quelea	<i>Quelea quelea</i>	40.00	30	7.41	2
488	Red-capped Lark	<i>Calandrella cinerea</i>	9.33	7	3.70	1
813	Red-collared Widowbird	<i>Euplectes ardens</i>	5.33	4	0.00	0
314	Red-eyed Dove	<i>Streptopelia semitorquata</i>	78.67	59	11.11	3
392	Red-faced Mousebird	<i>Urocolius indicus</i>	52.00	39	7.41	2
120	Red-footed Falcon	<i>Falco vespertinus</i>	2.67	2	0.00	0
820	Red-headed Finch	<i>Amadina erythrocephala</i>	30.67	23	0.00	0
940	Rock Dove	<i>Columba livia</i>	18.67	14	0.00	0
123	Rock Kestrel	<i>Falco rupicolus</i>	n/a			
506	Rock Martin	<i>Ptyonoprogne fuligula</i>	6.67	5	3.70	1
458	Rufous-naped Lark	<i>Mirafra africana</i>	36.00	27	3.70	1
460	Sabota Lark	<i>Calendulauda sabota</i>	8.00	6	3.70	1
789	Scaly-feathered Weaver	<i>Sporopipes squamifrons</i>	28.00	21	0.00	0
105	Secretarybird	<i>Sagittarius serpentarius</i>	2.67	2	0.00	0
847	Shaft-tailed Whydah	<i>Vidua regia</i>	1.33	1	3.70	1
504	South African Cliff Swallow	<i>Petrochelidon spilodera</i>	30.67	23	0.00	0
707	Southern Fiscal	<i>Lanius collaris</i>	72.00	54	14.81	4
709	Southern Boubou	<i>Laniarius ferrugineus</i>	1.33	1	0.00	0
4142	Southern Grey-headed Sparrow	<i>Passer diffusus</i>	26.67	20	3.70	1
803	Southern Masked Weaver	<i>Ploceus velatus</i>	78.67	59	3.70	1

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
808	Southern Red Bishop	<i>Euplectes orix</i>	64.00	48	3.70	1
390	Speckled Mousebird	<i>Collus striatus</i>	16.00	12	3.70	1
311	Speckled Pigeon	<i>Columba guinea</i>	69.33	52	11.11	3
474	Spike-heeled Lark	<i>Chersomanes albofasciata</i>	28.00	21	0.00	0
368	Spotted Eagle-Owl	<i>Bubo africanus</i>	2.67	2	0.00	0
654	Spotted Flycatcher	<i>Muscicapa striata</i>	17.33	13	0.00	0
275	Spotted Thick-knee	<i>Burhinus capensis</i>	6.67	5	0.00	0
88	Spur-winged Goose	<i>Plectropterus gambensis</i>	10.67	8	0.00	0
867	Streaky-headed Seedeater	<i>Crithagra gularis</i>	4.00	3	0.00	0
185	Swainson's Spurfowl	<i>Pternistis swainsonii</i>	40.00	30	0.00	0
649	Tawny-flanked Prinia	<i>Prinia subflava</i>	6.67	5	0.00	0
277	Temminck's Courser	<i>Cursorius temminckii</i>	n/a			
736	Violet-backed Starling	<i>Cinnyricinclus leucogaster</i>	1.33	1	0.00	0
840	Violet-eared Waxbill	<i>Granatina granatina</i>	5.33	4	0.00	0
735	Wattled Starling	<i>Creatophora cinerea</i>	49.33	37	0.00	0
359	Western Barn Owl	<i>Tyto alba</i>	2.67	2	0.00	0
61	Western Cattle Egret	<i>Bubulcus ibis</i>	49.33	37	3.70	1
391	White-backed Mousebird	<i>Collus colius</i>	46.67	35	3.70	1
107	White-backed Vulture	<i>Gyps africanus</i>	13.33	10	0.00	0
763	White-bellied Sunbird	<i>Cinnyris talatala</i>	13.33	10	0.00	0
780	White-browed Sparrow-Weaver	<i>Plocepasser mahali</i>	69.33	52	18.52	5
588	White-browed Scrub Robin	<i>Cercotrichas leucophrys</i>	1.33	1	0.00	0
409	White-fronted Bee-eater	<i>Merops bullockoides</i>	10.67	8	0.00	0
383	White-rumped Swift	<i>Apus caffer</i>	26.67	20	0.00	0
582	White-throated Robin-Chat	<i>Cossypha humeralis</i>	n/a			

#	Common Name	Scientific Name	SABAP2 Reporting Rate			
			Full Protocol (%)	Number of cards	Ad hoc Protocol (%)	Number of cards
814	White-winged Widowbird	<i>Euplectes albonotatus</i>	20.00	15	3.70	1
599	Willow Warbler	<i>Phylloscopus trochilus</i>	10.67	8	3.70	1
866	Yellow Canary	<i>Crithagra flaviventris</i>	66.67	50	7.41	2
96	Yellow-billed Duck	<i>Anas undulata</i>	34.67	26	0.00	0
129	Yellow-billed Kite	<i>Milvus aegyptius</i>	12.00	9	7.41	2
812	Yellow-crowned Bishop	<i>Euplectes afer</i>	9.33	7	0.00	0
859	Yellow-fronted Canary	<i>Crithagra mozambica</i>	2.67	2	0.00	0
629	Zitting Cisticola	<i>Cisticola juncidis</i>	33.33	25	0.00	0